Unintended Bias in Health Care Strategies for Providing More Equitable Care

BY MICHELLE VAN RYN, PHD, MPH

Research shows that unintentional bias on the part of physicians can influence the way they treat patients from certain racial and ethnic groups. Most physicians are unaware that they hold such biases, which can unknowingly contribute to inequalities in health care delivery. This article explains why a person's thoughts and behaviors may not align, and provides strategies for preventing implicit biases from interfering with patient care.

ver the past two decades, hundreds of studies have documented widespread inequalities in medical care. Although the reason for unequal care is multifaceted, physicians' behavior and decisions are known contributors.¹⁻⁸ Physicians' clinical decisions and the way they use of guidelines and evidence-based practices have been shown to contribute to disparities in:

- Care for cardiovascular risk factors ranging from hypertension^{9,10} to sleep disorders^{11,12}
- Treatment of symptoms associated with coronary artery disease and severe cardiac events^{6,13,14}
- Cancer screening, prevention, treatment and symptom management¹⁵
- Pediatric care, including asthma treatment^{16,17}
- Assessment, treatment and referral for mental health services.¹⁸

Disparities also have been shown between patients of different ages, racial and ethic groups, and genders receiving pain control.¹⁹⁻²² The questions physicians ask during patient interviews and the tests they order can contribute to such disparities in care.²³⁻²⁸

Physicians often find it difficult to accept that unconscious biases may affect the care they provide because the notion is so inconsistent with their explicit (conscious) attitudes, motivations and intentions. Most physicians have genuinely egalitarian conscious beliefs and want to provide excellent care to all of their patients.²⁹ The apparent contradiction between what they consciously believe and what research shows they actually do can cause considerable cognitive dissonance-the uncomfortable feeling people get when holding two conflicting ideas simultaneously. Cognitive dissonance is so uncomfortable that we will go to great lengths to resolve it, often discounting or ignoring evidence that supports the lesser-preferred of our beliefs. When physicians reject evidence of unconscious bias, they miss an opportunity to improve the quality of care they provide, thus potentially perpetuating the delivery of unequal care. This article is intended to help physicians resolve the cognitive dissonance they may feel related to disparities in health care by 1) explaining why-despite their best intentions-they might behave in ways inconsistent with their conscious beliefs and 2) providing strategies to prevent deep-seated biases from negatively affecting the care they provide.

Why Our Thoughts and Behaviors May Not Align

The reason physicians may be consciously well-intentioned yet behave in biased ways is rooted in the fact that we do not think the way we think we think. The vast majority of scientists studying the mind agree that humans have at least two separate information-processing systems that operate simultaneously. Daniel Kahneman, the Nobel Prize-winning author of *Thinking, Fast and Slow*, dubbed these simply as

System 1 and System 2.³⁰ We are primarily aware of System 2, which involves deliberative, reasoned, conscious and effortful thought. In contrast, System 1 often operates outside of our awareness, helping us navigate the millions of bits of information to which we are exposed at any one time by providing an unconscious framework for interpreting incoming information. For example, most people in North America will automatically make the association between an apple and food. Furthermore, when we see an apple, we will automatically draw on stored information about apples, avoiding the need to dissect and study every apple we encounter. Although most of us believe System 2-conscious and reasoned thought-guides our behavior and understanding of the world, Kahneman points out that, "System 1 is really the one that is the more influential ... it is steering System 2 to a very large extent."30

System 1 also guides us through our social interactions—and can sometimes lead us astray. For example, if white and Asian doctors are repeatedly exposed to blacks portrayed as criminals, violent or in other negative ways on television or in film, they may automatically and unconsciously associate black patients with threat and undesirable behavior. These unconscious expectations and attitudes, referred to as implicit biases, represent the "thumbprint of the culture on our minds"³¹ and, as such, they can be very different from our conscious attitudes and motives.

How Bias Manifests in the Clinic

Implicit biases have the potential to influence us in unintentional but powerful ways. Implicit racial bias has been shown to influence physicians' clinical decisionmaking^{3,32} in regard to patient referrals for thrombolysis³ and post-operative pain control for children.³³ Furthermore, implicit biases have been shown to have complex and subtle effects on physicianpatient interactions.³⁴ For example, physicians' level of implicit racial bias against blacks, as assessed by the Implicit Associations Test, have been found to be inversely associated with patient-centered behavior,^{8,35,36} visit length,³⁵ warmth,⁴ and positively associated with rapidity of speech35,37 and verbal dominance during the encounter.³⁸ Studies showed black patients reported less respect for, confidence in, and trust in the advice of medical professionals who scored higher in implicit bias.^{34,35} This distrust predicted lower levels of adherence to the physician's recommendations,³⁹ a finding consistent with other evidence that patients' perceptions of being judged, negatively perceived, stigmatized or discriminated against predict patient adherence 40-44 and likelihood of seeking followup or preventive care.^{38,45-52}

It is important to bear in mind that implicit bias is not unique to physicians or the health care industry. Examples of the pernicious effects of implicit racial and other biases exist in every sector of our society. For example, fictitious applicants with identical resumes responding to 1,300 want ads got 50% more call-backs when they used a "white-sounding" name versus a "black-sounding" name.53 Another found female musicians were significantly less likely than male musicians to be hired for orchestras during open auditions, but as or more likely to be hired when they auditioned from behind a curtain.⁵⁴ In yet another study, faculty members (both male and female) reviewing applications for a student lab manager position that were identical except for gender viewed the male applicants as more competent than the female applicants. They also were more likely to hire and mentor

a male student than a female student and offer him a higher starting salary.55

Strategies for Providing More Equitable Care

Although our implicit biases can cause us to behave in ways that are inconsistent with our explicit motives, values and beliefs, they do not have to. There are strategies that can increase our likelihood of seeing patients in terms of their unique individual characteristics, as opposed to those of a social or cultural group of which they are a member. In a recent issue of Minnesota Medicine, editor in chief Charles Meyer, MD, described the challenge: "Equity in the exam room means treating each patient as if they were your most important patient, regardless of gender, sexual orientation, race, ethnicity or personal appearance." 56

The massive body of evidence demonstrating the negative impact of implicit bias has prompted a number of additional studies identifying factors that can minimize it. The following are recommendations from those studies that have the strongest supporting evidence.

1. Put yourself in your patients' shoes. Numerous studies have found that perspective-taking reduces bias and inhibits the activation of unconscious stereotypes and prejudices.^{34,38,57,58} Perspective-taking refers to imagining yourself in the other person's position; seeing things through his or her eyes. It is the cognitive component of empathy, and it can be learned and cultivated with practice. In addition to its documented benefits for reducing bias and stereotypes, provider empathy has been associated with increased patient satisfaction, adherence to physicians' recommendations, self-efficacy and perceptions of control; less emotional distress; and better outcomes. 59,60 Some physicians have highly developed perspective-taking skills. But even those who do may not routinely apply them during clinical encounters. Through daily practice with family, friends and colleagues these skills can improve over time and their use will become more automatic.

Steps you can take:

- Imagine yourself in the other person's shoes. Think of it as walking in their world or seeing the world through their eyes.
- Check in with your patient by saying something like: "I am wondering how I might see the situation if I were looking through your eyes ... " or "I was imagining being in your shoes, and it occurred to me that I might (feel/think/be) Am I close?"
- Read essays, narratives and fiction that provide the point of view of others who differ from you in terms of culture, race/ethnicity, socioeconomic status or another characteristic.

2. Build partnerships with your patients. Cultivate a sense that you and your patient (and perhaps his or her family) are on the same team, working toward shared goals. Being in partnership with patients creates a sense of a common ingroup identity and reduces the likelihood of being "hijacked" by implicit biases.⁶¹⁻⁶⁴ Research has shown that we like, trust and are more motivated to help people in our "in group"-those we believe to be like us.⁶⁵⁻⁶⁷ We tend to attribute the problematic behavior of members of our in-group to situational factors (eg, he was confused by the instructions), whereas we tend to attribute such behaviors among those who are not members of our in-group to an individual's intelligence or personality. For example, a white physician may describe an African-American patient who failed to take her medications as instructed as "nonadherent," yet that same physician might say a white patient who didn't follow her instructions for taking the medication "forgot the timing" or "needs additional instruction." Such attributions may cumulatively affect future encounters with those patients. Thus, the value of developing a partnership with patients and creating a sense of the patient being a member of ones' in-group can reduce categorization and associated implicit bias.67-69 Partnership-building also promotes rapport and patient trust, potentially improving adherence and outcomes.

Steps you can take:

- Use the terms "we" and "us" instead of "I" and "you" to make it feel as if you're all members of the same team.⁶² For example, instead of "*I* am going to order X test," try "*We* should probably use X test so we can find out…" or "*Let's* use X test." Instead of "*I* am going to prescribe Y" try "*Our* best course of action might be to try Y." Rather than say "If *you* have Y side effects..." try "If *we* find that Y side effects are a problem..."
- Focus on your common goals. It can help to articulate them by saying: "It seems as if our most important goal is to... (reduce symptoms, cure X, prevent Y, etc.)." This also helps prevent misunderstandings by allowing the patient to clarify or discuss them.
- Listen attentively and responsively, invite patients to participate in clinical decision-making, focus on the patient's strengths (and help that patient focus on their strengths), validate the patient's perspectives and concerns, and respect and honor their values.

3. Take care of yourself-protect your mental resources. Physicians and other health care providers are notorious for caring for others at the expense of their own well-being. However, converging lines of research suggest that self-care and emotional regulation skills are crucial to providing high-quality, unbiased care. Studies have shown that when people have sufficient motivation, resources, information, time and awareness to be mindful, their judgement, behavior and decision-making are much less likely to be undermined by implicit biases.⁶⁹⁻⁷³ However, when illness, fatigue, stress, anxiety or competing demands command more of their mental resources, their cognitive processing capacity may be compromised, allowing implicit biases and attitudes to hijack perceptions, expectations and evaluations of patients. Unfortunately, competing demands, distractions, heavy workloads and time pressure-all of which can increase stress and fatigue and decrease cognitive capacityare all too common in clinical settings.⁷⁴

Steps you can take:

- Assess your practice for unnecessary cognitive demands. This may mean addressing such things as scheduling, noise levels, inadequate training, poor supervision and clinic or facility overcrowding.⁷⁵
- Allow adequate time per patient and between patients, establish routines and make sure your clinic has sufficient staffing. ⁷⁴
- Do things to protect your mental energy, such as getting sufficient sleep, finding ways to reduce stress and taking mental breaks throughout the day to refocus on being present with your patients.

4. Be positive. Research suggests that physicians who have positive emotions during the clinical encounter are more likely to see their patients as unique individuals and/or part of their in-group, and less likely to categorize them in terms of their race, ethnicity or culture. ^{67,76}

Steps you can take:

- Strengthen or add practices associated with positive mental health such as mindfulness-based stress reduction, regular physical exercise, engagement in a pleasant hobby or sport, and time with friends and family. Scheduled solitude, if you are a person who benefits from time alone.
- Learn and use strategies for rapidly shifting negative emotions, especially those caused by stress or anxiety. Examples include abdominal breathing techniques, progressive muscle relaxation, mindfulness, and/or focusing for a moment on something you appreciate or for which you feel grateful.

5. Counter negative stereotypes by exposing yourself to positive images. Our implicit biases reflect ideas repeated in the larger society. One way to reduce our own biases is to expose ourselves to images that differ from what we commonly see. Studies have shown that exposure to admired African Americans and to images of African Americans in positive settings reduced negative implicit bias on the part of whites.^{77,78} Steps you can take:

- Seek out entertainment that portrays racial and ethnic minorities in positive roles; women as likeable, competent leaders; obese people as active and intelligent; and elderly people as intellectually sharp and productive.
- Display artwork that portrays members of various groups in a positive light. Having artwork in waiting rooms, hallways and exam rooms that counters stereotypes may both reduce negative bias and make diverse patients feel valued. Even engaging in mental imagery that involves counter-stereotypical representations has shown benefit.⁷⁹
- Bring groups of diverse people together to work toward a common goal. A metaanalysis of 515 studies concluded that intergroup contact typically reduces intergroup bias and intergroup anxiety.⁸⁰

Conclusion

Many people, physicians included, believe that the problem of implicit bias only applies to other people, even though research suggests that almost all of us have negative implicit attitudes toward people from various groups. But these implicit biases do not have to control our behavior. By engaging in self-awareness, being mindful, regulating our emotions, routinely practicing perspective-taking, building relationships with people in other groups, practicing self-care and protecting our mental energy, we can go a long way toward ensuring that our behavior toward others reflects our true values, goals and motives. MM

Michelle van Ryn is director of the research program on equity and inclusion in health care at Mayo Clinic and executive director of Partners in Equity and Inclusion.

REFERENCES

1. Cooper LA. Overcoming Healthcare Disparities: The Role of Patient-Centered Care. College of Public Health and Health Professions Celebrating 50 Years. University of Florida 2008.

2. Dovidio JF, Penner LA, Albrecht TL, Norton WE, Gaertner SL, Shelton JN. Disparities and distrust: the implications of psychological processes for understanding racial disparities in health and health care. Soc Sci Med.2008;67(3):478-86.

3. Green AR, Carney DR, Pallin DJ, et al. Implicit bias among physicians and its prediction of thrombolysis decisions for black and white patients. J Gen Intern Med. 2007;22(9):1231-8.

4. Penner LA, Dovidio JF, West TV, et al. Aversive racism and medical interactions with black patients: a field study. J Exp Soc Psychol. 2010;46(2):436-40.

5. Smedley BD, Stith AY, Nelson AR (eds). Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care. National Academy Press; Washington, DC, 2003.

6. van Ryn M, Burgess D, Malat J, Griffin J. Physicians' perceptions of patients' social and behavioral characteristics and race disparities in treatment recommendations for men with coronary artery disease. Am J Public Health. 2006;96(2):351-7.

7. van Ryn M, Fu SS. Paved with good intentions: do public health and human service providers contribute to racial/ethnic disparities in health? Am J Public Health. 2003;93(2):248-55.

8. Cooper LA, Roter DL, Carson KA, et al. The associations of clinicians' implicit attitudes about race with medical visit communication and patient ratings of interpersonal care. Am J Public Health. 2012;102(5):979-87.

9. Eapen ZJ, Liang L, Shubrook JH, et al. Current quality of cardiovascular prevention for Million Hearts: An analysis of 147,038 outpatients from The Guideline Advantage. Am Heart J. 2014;168(3):398-404.

10. Will JC, Nwaise IA, Schieb L, Zhong Y. Geographic and racial patterns of preventable hospitalizations for hypertension: Medicare beneficiaries, 2004-2009. Public Health Rep. 2014;129(1):8-18.

11. Reducing Health Disparities: The Role of Sleep Deficiency and Sleep Disorders Bethesda, MD: National Heart, Lung, and Blood Institute; December 2011.

12. Kingsbury JH, Buxton OM, Emmons KM. Sleep and its relationship to racial and ethnic disparities in cardiovascular disease. Curr Cardiovasc Risk Rep. 2013;7(5).

13. Bradley EH, Herrin J, Wang Y, et al. Racial and ethnic differences in time to acute reperfusion therapy for patients hospitalized with myocardial infarction. JAMA. 2004;292(13):1563-72.

14. Hannan EL, van Ryn M, Burke J, et al. Access to coronary artery bypass surgery by race/ethnicity and gender among patients who are appropriate for surgery. Med Care. 1999;37(1):68-77.

15. Shavers VL, Brown ML. Racial and ethnic disparities in the receipt of cancer treatment. J Natl Cancer Inst. 2002;94(5):334-57.

16. National Heart, Lung and Blood Institute. Reducing Asthma Disparities. Available at: www. nhlbi.nih.gov/health-pro/resources/lung/naci/discover/ disparities.htm. Accessed February 9, 2016.

17. Partners Putting Guidelines Into Action: Expert Panel Report 3 -- Guidelines for the Diagnosis and Management of Asthma. National Heart, Lung, and Blood Institute, National Institutes of Health, Department of Health and Human Services. December 2008.

18. Meyer OL, Saw A, Cho YI, Fancher TL. Disparities in assessment, treatment, and recommendations for specialty mental health care: patient reports of medical provider behavior. Health Serv Res. 2015;50(3):750-67.

19. Hwang U, Belland LK, Handel DA, et al. Is all pain is treated equally? A multicenter evaluation of acute pain care by age. Pain. 2014;155(12):2568-74.

20. Todd KH. Influence of ethnicity on emergency department pain management. Emerg Med (Fremantle). 2001;13(3):274-8.

21. Todd KH, Deaton C, D'Adamo AP, Goe L. Ethnicity and analgesic practice. Ann Emerg Med. 2000;35(1):11-6.

22. Hampton SB, Cavalier J, Langford R. The influence of race and gender on pain management: a systematic literature review. Pain Manag Nurs. 2015;16(6):968-77.

23. Eggly S, Harper FW, Penner LA, Gleason MJ, Foster T, Albrecht TL. Variation in question asking during cancer clinical interactions: A potential source of disparities in access to information. Patient Educ Couns. 2011;82(1):63-8

24. Durkin MS, Maenner MJ, Meaney FJ, et al. Socioeconomic inequality in the prevalence of autism spectrum disorder: evidence from a U.S. cross-sectional study. PLoS One. 2010;5(7):e11551.

25. Miller TW, Nigg JT, Miller RL. Attention deficit hyperactivity disorder in African American children: what can be concluded from the past ten years? Clin Psychol Rev. 2009;29(1):77-86.

26. Alegria M, Nakash O, Lapatin S, et al. How missing information in diagnosis can lead to disparities in the clinical encounter. J Public Health Manag Pract. 2008;14 Suppl:S26-35.

27. Wiehe SE, Rosenman MB, Wang J, Fortenberry JD. Disparities in chlamydia testing among young women with sexually transmitted infection symptoms. Sex Transm Dis. 2010 37(12):751-5.

28. Quinn K, Shalowitz MU, Berry CA, Mijanovich T, Wolf RL. Racial and ethnic disparities in diagnosed and possible undiagnosed asthma among publicschool children in Chicago. Am J Public Health. 2006;96(9):1599-603.

29. Epstein RA. Disparities and discrimination in health care coverage: a critique of the Institute of Medicine study. Perspect Biol Med. 2005;48(1 Suppl):S26-41.

30. Kahneman DT. Thinking, fast and slow. New York: Farrar, Straus and Giroux; 2011.

31. Banaji MR, Greenwald A. Blindspot: Hidden Biases of Good People. Delacorte Press; 2013.

32. Sabin JA, Marini M, Nosek BA. Implicit and explicit anti-fat bias among a large sample of medical doctors by BMI, race/ethnicity and gender. PLoS One. 2012;7(11):e48448.

33. Sabin JA, Greenwald AG. The influence of implicit bias on treatment recommendations for 4 common pediatric conditions: pain, urinary tract infection, attention deficit hyperactivity disorder, and asthma. Am J Public Health. 2012;102(5):988-95.

34. Dovidio JF, Penner LA, Albrecht TL, Norton WE, Gaertner SL, Shelton JN. Disparities and distrust: the implications of psychological processes for understanding racial disparities in health and health care. Soc Sci Med. 2008;67(3):478-86.

35. Cooper LA, Roter DL, Carson KA, et al. The associations of clinicians' implicit attitudes about race with medical visit communication and patient ratings of interpersonal care. Am J Public Health. 2012;102(5):979-87.

36. Blair IV, Havranek EP, Price DW, et al. Assessment of biases against Latinos and African Americans among primary care providers and community members. Am J Public Health. 2013;103(1):92-8.

37. Blair IV, Steiner JF, Fairclough DL, et al. Clinicians' implicit ethnic/racial bias and perceptions of care among black and latino patients. Ann Fam Med. J 2013;11(1):43-52.

38. Hagiwara N, Penner LA, Gonzalez R, et al. Racial attitudes, physician-patient talk time ratio, and adherence in racially discordant medical interactions. Soc Sci Med. 2013;87:123-31.

39. Penner LA, Dovidio JF, Edmondson D, et al. The experience of discrimination and black-white health disparities in medical care. J Black Psychol. 2009;35(2). **40**. Martin KD, Roter DL, Beach MC, Carson KA, Cooper LA. Physician communication behaviors and trust among black and white patients with hypertension. Med Care. 2013;51(2):151-7.

41. Cuffee YL, Hargraves JL, Rosal M, et al. Reported racial discrimination, trust in physicians, and medication adherence among inner-city African Americans with hypertension. Am J Public Health. 2013;103(11):e55-62.

42. Omosanya OE, Elegbede OT, Agboola SM, Isinkaye AO, Omopariola OA. Effects of stigmatization/discrimination on antiretroviral therapy adherence among HIV-infected patients in a rural tertiary medical center in Nigeria. J Int Assoc Provid AIDS Care. 2014;13(3):260-3.

43. Kronish IM, Diefenbach MA, Edmondson DE, Phillips LA, Fei K, Horowitz CR. Key barriers to medication adherence in survivors of strokes and transient ischemic attacks. J Gen Intern Med. 2013;28(5):675-82.

44. Gudzune KA, Bennett WL, Cooper LA, Bleich SN. Perceived judgment about weight can negatively influence weight loss: A cross-sectional study of overweight and obese patients. Prev Med. 2014;62:103-7.

45. Amy NK, Aalborg A, Lyons P, Keranen L. Barriers to routine gynecological cancer screening for White and African-American obese women. Int J Obes (Lond). 2006;30(1):147-55.

46. Anderson DA, Wadden TA. Bariatric surgery patients' views of their physicians' weight-related attitudes and practices. Obes Res. 2004;12(10):1587-95.

47. Aldrich T, Hackley B. The impact of obesity on gynecologic cancer screening: an integrative literature review. J Midwifery Womens Health. 2010;55(4):344-56.

48. Drury CA, Louis M. Exploring the association between body weight, stigma of obesity, and health care avoidance. J Am Acad Nurse Pract. 2002;14(12):554-61.

49. Jones PR, Taylor DM, Dampeer-Moore J, et al. Health-related stereotype threat predicts health services delays among blacks. Race Soc Probl. 2013;5(2):121-36.

50. Levine CS, Ambady N. The role of non-verbal behaviour in racial disparities in health care: implications and solutions. Med Educ. 2013;47(9):867-76.

51. Eggly S, Barton E, Winckles A, Penner LA, Albrecht TL. A disparity of words: racial differences in oncologist-patient communication about clinical trials. Health Expect. 2013;18(5):1316:26.

52. Dovidio JF, Fiske ST. Under the radar: how unexamined biases in decision-making processes in clinical interactions can contribute to health care disparities. Am J Public Health. 2012;102(5):945-52.

53. Bertrand M, Mullainathan S. Are Emily and Greg more employable than Lakisha and Jamal? A field experiment on labor market discrimination. Am Econ Rev. 2004;94(4):991-1013.

54. Goldin C, Rouse C. Orchestrating impartiality: the impact of "blind" auditions on female musicians. Am Econ Rev. 2000;90(4):715-41.

55. Moss-Racusin CA, Dovidio JF, Brescoll VL, Graham MJ, Handelsman J. Science faculty's subtle gender biases favor male students. Proc Natl Acad Sci U S A. 2012;109(41):16474-9.

56. Meyer C. Beneath the surface. Minn Med. 2015;98(10):4.

57. Galinsky AD, Moskowitz GB. Perspective-taking: decreasing stereotype expression, stereotype accessibility, and in-group favoritism. J Pers Soc Psychol. 2000;78(4):708-24.

(continued on page 46)

e-prescribing and EHR vendors have not yet incorporated ePA functionality. The Table summarizes the status of ePA functionality of the five most commonly used EHRs in Minnesota clinics at the time of this writing. Other EHR vendors that also lack ePA functionality at the time of this writing are AmazingCharts, athenahealth, e-MDs, GE Healthcare, Greenway Health, McKesson and Meditech. Physician practices are encouraged to contact their vendors directly to learn whether their EHR has ePA functionality, and if not when it's expected to be available and if any version updates will be needed. If they do not have access to ePA through their EHR, physicians may be able to achieve compliance with the state mandate using web-based services such as CoverMyMeds and PARx.

Another challenge has been slow adoption of ePA by payers and PBMs. Some national payers may not yet have ePA capabilities. Several large companies including Aetna, Cigna and UnitedHealthcare appear to be on track even though they have limited market penetration in Minnesota. Among payers that do not have such capabilities at the time of this writing are the Minnesota Department of Human Services (Medical Assistance and MinnesotaCare) and HealthPartners. The state has indicated that it will not be able to accept electronic requests for medication prior authorizations until at least April 1, 2016. HealthPartners has not yet provided a go-live date.

Conclusion

Full adoption of ePA for medications offers enormous potential for reducing the time and cost associated with negotiating variable insurance requirements and labor-intensive administrative processes. Physicians and patients may still experience challenges accessing formularies and obtaining coverage for medications, but ePA should facilitate communication between prescribers and payers and improve the prior authorization process for all parties. MM Janet Silversmith is director of health policy and member services for the Minnesota Medical Association. Laura Topor is president of Granada Health, Inc. and a past board member of the National Council of Prescription Drug Programs.

REFERENCES

 Morra D, Nicholson S, Levinson W, Gans D, Hammons T, Casalino L. US physician practices versus Canadians: spending nearly four times as much money interacting with payers. Health Aff (Millwood). 2011;30(8):1443-50.

2. Soumerai S. Benefits and risks of increasing restrictions on access to costly drugs in Medicaid. Health Aff (Millwood). 2004;23(1):135-46.

Minnesota Statute § 62J.497

4. Minnesota Statute § 62J.497 Subd. 5.

(continued from page 43)

58. Finlay KA, Stephan WG. Improving intergroup relations: The effects of empathy on racial attitudes. J Applied Soc Psychol. 2000;30(8):1720-37.

59. Dwamena F, Holmes-Rovner M, Gaulden CM, et al. Interventions for providers to promote a patient-centred approach in clinical consultations. Cochrane Database Syst Rev. 2012;12:CD003267.

60. Kelley JM, Kraft-Todd G, Schapira L, Kossowsky J, Riess H. The influence of the patient-clinician relationship on healthcare outcomes: a systematic review and meta-analysis of randomized controlled trials. PLoS One. 2014;9(4):e94207.

61. Penner LA, Hagiwara N, Eggly S, Gaertner SL, Albrecht TL, Dovidio JF. Racial healthcare disparities: a social psychological analysis. Eur Rev Soc Psychol. 2013;24(1):70-122.

62. Dovidio JF, Gaertner SL, Saguy T. Commonality and the complexity of "we": social attitudes and social change. Pers Soc Psychol Rev. 2009;13(1):3-20.

63. Gomez A, Dovidio JF, Huici C, Gaertner SL, Cuadrado I. The other side of we: when outgroup members express common identity. Pers Soc Psychol Bull. 2008;34(12):1613-26.

64. Dovidio JF, Penner LA, Albrecht TL, Norton WE, Gaertner SL, Shelton JN. Disparities and distrust: the implications of psychological processes for understanding racial disparities in health and health care. Soc Sci Med. 2008;67(3):478-86.

65. Fiske ST. Varieties of (de) humanization: divided by competition and status. Nebr Symp Motiv. 2013;60:53-71.

66. Cikara M, Fiske ST. Bounded empathy: neural

responses to outgroup targets' (mis)fortunes. J Cogn Neurosci. 2011;23(12):3791-3803.

67. Dovidio JF, Gaertner SL, Validzic A. Intergroup bias: status, differentiation, and a common in-group identity. J Pers Soc Psychol. 1998;75(1):109-20.

68. Penner LA, Gaertner S, Dovidio JF, et al. A social psychological approach to improving the outcomes of racially discordant medical interactions. J Gen Intern Med. 2013;28(9):1143-9.

69. Gaertner SL, Dovidio JF, Rust MC, et al. Reducing intergroup bias: elements of intergroup cooperation. J Pers Soc Psychol. 1999;76(3):388-402.

70. Kang Y, Gray JR, Dovidio JF. The nondiscriminating heart: lovingkindness meditation training decreases implicit intergroup bias. J Experiment Psychol Gen. 2014;143(3):1306-13.

71. Lebowitz MS, Dovidio JF. Implications of emotion regulation strategies for emphathic concern, social attitudes and helping behavior. Emotion. 2015;15(2):187-94.

72. Devine PG, Forscher PS, Austin AJ, Cox WT. Long-term reduction in implicit race bias: A prejudice habit-breaking intervention. J Exp Soc Psychol. 2012;48(6):1267-78.

73. Stewart BD, Payke BK. Bringing automatic stereotyping under control: implementation intentions as efficient means of thought control. Pers Soc Psychol Bull. 2008;34(10):1332-45.

74. Burgess DJ. Are providers more likely to contribute to healthcare disparities under high levels of cognitive load? How features of the healthcare setting may lead to biases in medical decision making. Med Decis Making. 2010;30(2):246-57.

75. Crockskerry P, Wears R. Safety errors in emergency medicine. In: Markovchick V, Pons P, eds. Emergency Medicine Secrets. 3rd ed. Philadelphia: Hanley and Belfus; 2002:29-37.

76. Johnson KJF, Barbara L. "We all look the same to me": positive emotions eliminate the own-race bias in face recognition. Psychol Sci. 2005;16:875-81.

77. Allen TJ, Sherman JW, Klauer KC. Social context and the self-regulation of implicit bias. Group Processes Intergroup Relations. 2010;13(4):137-49.

78. Lai CK, Marini M, Lehr SA, et al. Reducing implicit racial preferences: I. A comparative investigation of 17 interventions. J Exp Psychol Gen. 2014;143(4):1765-85.

79. Blair IV, Ma JE, Lenton AP. Imagining stereotypes away: the moderation of implicit stereotypes through mental imagery. J Pers Soc Psychol. 2001;81(5):828-41.

80. Pettigrew T, Tropp L. How does intergroup contact reduce prejudice? Meta-analytic tests of three mediators. Euro J Soc Psychol. 2008;38(6):922-34.